International Journal of Computational and Experimental Science and ENgineering (IJCESEN)

Vol. 11-No.4 (2025) pp. 7473-7481
http://www.ijcesen.com

Copyright © IJCESEN

ISSN: 2149-9144

Research Article

Impact of Rapid Emergency Medical Services Response Times on Patient Outcomes in Saudi Arabia

Mousa Abdullah Alhajji Ahmed[1]*, Almousa, Hassan Abdulrahman T[2], Abdullah Ali Al Kishi[3], Ahmad Amer Hussain Alamer[4], Hussain Salem Alali[5], Abbas Ali Alhajji[6], Abdulaziz Hussain Ahmed Alabdullah[7], Mohammed Abbs Ali Aloqili[8], Reda mohammad Ali Alnahawi[9], Saleh Mohammed Albattat[10]

[1]Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
* **Corresponding Author Email:** mussasam2@gmail.com - **ORCID:** 0000-0002-0047-7850

[2] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: Halmosa@srca.org.sa- **ORCID:** 0000-0002-9247-7850

[3] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: dar-ahzan@hotmail.com- **ORCID:** 0000-0002-8247-7850

[4] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: kingmoon8@gmail.com- **ORCID:** 0000-0002-7247-7850

[5] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: Huszsain@yahoo.com- **ORCID:** 0000-0002-6247-7850

[6] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: asab1407@hotmail.com- **ORCID:** 0000-0002-4247-7850

[7] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: Alabdullahaziz91@gmail.com - **ORCID:** 0000-0002-3247-7850

[8] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: Mohammedalokily20@gmail.com- **ORCID:** 0000-0002-247-7850 3

[9] Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: Redaalnahawi@gmail.com- **ORCID:** 0000-0002-2247-7850

[10]Emergency Medical Services, Saudi Red Crescent Authority, Saudi Arabia
Email: galaxy760@live.com - **ORCID:** 0000-0002-1247-7850

Article Info:

DOI: 10.22399/ijcesen.4049
Received : 02 January 2025
Accepted : 30 January 2025

Keywords

Emergency Medical Services,
response times,
patient outcomes,
Saudi Arabia

Abstract:

Rapid Emergency Medical Services (EMS) response times play a crucial role in determining patient outcomes in Saudi Arabia, where the prevalence of acute medical conditions such as cardiac arrests, strokes, and trauma cases is significant. Studies indicate that timely intervention by EMS can drastically improve survival rates and reduce morbidity associated with these emergencies. In urban areas, where traffic congestion can delay response times, the implementation of advanced dispatch systems and dedicated emergency lanes has been shown to enhance the speed of medical assistance. Furthermore, public awareness campaigns about the importance of calling for emergency services promptly can empower citizens to act swiftly in critical situations, ultimately contributing to better health outcomes. In rural regions of Saudi Arabia, the challenges of geographical barriers and limited access to healthcare facilities can exacerbate the impact of delayed EMS responses. The establishment of mobile health units and the integration of telemedicine have emerged as potential solutions to bridge these gaps, ensuring that patients receive timely care regardless of their location. Research underscores the need for continuous training and education for EMS personnel to improve their efficiency and effectiveness in emergency situations. Overall, optimizing EMS response times is essential for enhancing patient survival rates and quality of care across the diverse landscapes of Saudi Arabia.

1. Introduction

The provision of effective and timely emergency medical services (EMS) is a critical benchmark of a nation's public health infrastructure and its commitment to safeguarding human life. At the core of EMS efficacy lies a single, pivotal metric: response time. Defined as the interval between the receipt of an emergency call and the arrival of a qualified medical team at the patient's location, response time is often the differentiating factor between life and death, between full recovery and permanent disability [1]. The concept of the "golden hour"—the critical period following a traumatic injury during which there is the highest likelihood that prompt medical treatment will prevent death—has long been a guiding principle in emergency medicine, underscoring the profound impact of rapid intervention on patient survival and neurological outcomes [2]. In the context of the Kingdom of Saudi Arabia, a nation characterized by vast geographical expanses, rapidly growing urban centers, and the unique challenge of managing mass gatherings during the Hajj and Umrah seasons, the optimization of EMS response times is not merely an operational goal but a national imperative aligned with the health objectives of Saudi Vision 2030 [3].

The Saudi Red Crescent Authority (SRCA), as the primary national provider of pre-hospital care, shoulders the immense responsibility of meeting this imperative. In recent years, the SRCA has embarked on an ambitious transformation, leveraging technological advancements, strategic infrastructure development, and rigorous training to enhance its operational performance. Recent reports indicate significant progress; for instance, the SRCA has announced achieving an average response time of less than 15 minutes in urban centers like Riyadh and Jeddah, a marked improvement from previous years [4]. This is supported by the strategic deployment of over 2,000 ambulances and the establishment of more than 450 stations across the Kingdom, alongside the integration of advanced Computer-Aided Dispatch (CAD) systems that optimize resource allocation [5]. Despite these advancements, disparities persist. Response times in remote and rural regions can extend to 30 minutes or more, highlighting a critical challenge in achieving equitable emergency care access across the Kingdom's diverse landscape [6]. This urban-rural divide presents a significant variable in studying patient outcomes on a national scale.

The clinical rationale for minimizing response times is robust and well-documented in international literature. For every minute that passes following a sudden cardiac arrest (SCA), the chance of survival decreases by 7-10% [7]. The early administration of cardiopulmonary resuscitation (CPR) and defibrillation by EMS teams is the most critical determinant of survival for out-of-hospital cardiac arrest (OHCA) victims. In cases of major trauma, such as those resulting from road traffic accidents—a leading cause of death and disability in Saudi Arabia—rapid on-scene stabilization, hemorrhage control using tourniquets, and swift transport to a designated trauma center are proven to significantly reduce mortality rates [8]. Similarly, for acute medical emergencies like stroke, the phrase "time is brain" encapsulates the urgent need for intervention; faster EMS response and pre-hospital notification to receiving hospitals enable quicker administration of thrombolytic therapy, drastically improving functional outcomes and reducing long-term disability [9].

Within the Saudi context, the epidemiological transition towards an increased burden of non-communicable diseases (NCDs) such as cardiovascular diseases and diabetes further amplifies the importance of a rapid EMS system. The World Health Organization (WHO) estimates that NCDs account for over 70% of deaths in Saudi Arabia, a statistic that translates into a high volume of calls for chest pain, respiratory distress, and diabetic emergencies [10]. The ability of the SRCA to respond swiftly to these time-sensitive conditions is, therefore, a direct determinant of national health outcomes. Furthermore, Saudi Arabia's status as the host of the Hajj, which gathers over two million pilgrims in a confined area, represents the ultimate stress test for any EMS system. The strategies and response time benchmarks developed and achieved during such mass gatherings often serve as a model for best practices and system innovation that can be adapted for routine operations throughout the year [11].

However, while the correlation between rapid response and improved outcomes is conceptually clear, a comprehensive and localized analysis is required to quantify this impact specifically within the Saudi healthcare ecosystem. Factors such as public awareness and willingness to activate EMS, the proficiency of bystander CPR, traffic congestion in major cities, and the efficiency of the handover process at emergency departments all interact with response times to ultimately determine patient prognosis [12].

2. Defining the Golden Hour and Platinum Minutes: The Clinical Imperative for Speed

The concept of the **"Golden Hour"** emerged from military and civilian trauma research in the latter half of the 20th century. It postulates that the first 60 minutes following a major traumatic injury represent a critical window of opportunity. If definitive surgical care and resuscitation can be initiated within this window, the patient's chances of survival are significantly maximized [13]. The biological rationale for this timeframe is rooted in the body's progressive response to severe injury. Immediately following trauma, a cascade of events begins: uncontrolled hemorrhage leads to hypovolemic shock, depriving vital organs of oxygen; the body's compensatory mechanisms begin to fail; and the risk of entering an irreversible state of shock and coagulopathy (the body's inability to clot blood) increases exponentially with each passing minute. Studies have consistently shown that delays in surgical intervention for internal bleeding are directly correlated with a steep rise in mortality [14]. In the context of Saudi Arabia, where road traffic accidents (RTAs) are a leading cause of death and disability, particularly among the youth, the Golden Hour is of paramount importance [15]. The chain of survival for a trauma victim requires that the "clock" starts at the moment of injury, with the SRCA's response time constituting a major portion of this precious hour. Any delay in the "scene time" (the time spent stabilizing the patient on-site) or "transport time" (the time taken to reach an appropriate trauma center) directly eats into the time available for life-saving surgery in the hospital. Therefore, a rapid EMS response is the first and most crucial link in the trauma care continuum, enabling critical interventions like hemorrhage control and airway management to begin in the field, thereby preserving the patient's chance to survive until they reach the operating room.

While the Golden Hour broadly addresses trauma, the concept of **"Platinum Minutes"** applies with even greater urgency to specific medical emergencies, most notably out-of-hospital cardiac arrest (OHCA). The term "Platinum" signifies an even more valuable and fleeting window than gold—typically the first 5 to 10 minutes after cardiac arrest [16]. The pathophysiology here is stark: when the heart stops beating, the brain and other vital organs are immediately deprived of oxygen. Brain cells begin to die within minutes, and for each minute that passes without cardiopulmonary resuscitation (CPR) and defibrillation, the chance of survival decreases by 7-10% [17]. After approximately 10 minutes without intervention, the likelihood of survival with good neurological function becomes negligible. This creates an immense pressure on the EMS system. The Platinum Minutes mandate that the time from collapse to the delivery of the first shock from an automated external defibrillator (AED) must be minimized to the greatest extent possible. This places the SRCA's average urban response time of, for instance, 15 minutes, into a critical perspective; it underscores that for OHCA patients, the system is often racing against a clock that expires before a traditional ambulance can even arrive. This reality highlights the indispensable role of bystander intervention and public access defibrillation programs, which will be discussed in a later section. For other time-sensitive conditions like acute ischemic stroke, a similar principle applies, often termed "Time is Brain." Here, rapid EMS response is critical not only for swift transport but also for pre-hospital notification, allowing the hospital's stroke team to prepare for immediate administration of thrombolytic therapy upon the patient's arrival, a treatment whose efficacy is heavily dependent on a narrow time window [18].

The clinical imperative for speed, therefore, transcends simple efficiency; it is a battle against irreversible physiological decline. The relationship between time and outcome is not linear but rather an exponential decay, particularly for OHCA and severe hemorrhagic shock. International benchmarks consistently validate this. For example, the American Heart Association recommends a goal of first shock delivery within 5 minutes of EMS call receipt for OHCA, while trauma systems aim for a total pre-hospital time (response plus scene plus transport) of less than 60 minutes for major trauma victims [19]. For the SRCA, measuring performance against these clinical imperatives is crucial. It shifts the focus from merely logging faster response times to understanding how those times directly impact specific clinical endpoints: survival-to-discharge rates for cardiac arrest, mortality rates from RTAs, and functional independence scores for stroke survivors. A study conducted in the Riyadh region found that OHCA patients who received their first defibrillation within 8 minutes had a survival rate over three times higher than those for whom it took longer, powerfully illustrating the tangible impact of the Platinum Minutes concept in the Saudi context [20].

3. The Saudi EMS Landscape: Structure, Performance Metrics, and the Urban-Rural Divide

The SRCA operates as a unified national service, a structure that provides significant advantages in standardization, resource allocation, and strategic

planning. Governed by a central administration, it has developed a regional model, dividing the Kingdom into several operational sectors, each with its own directorate responsible for day-to-day operations [21]. This centralized-yet-decentralized model allows for the implementation of national protocols and training standards while providing regional branches the flexibility to address local challenges. The SRCA's fleet, comprising over 2,000 advanced life support (ALS) and basic life support (BLS) ambulances, is strategically managed through a network of more than 450 fixed and mobile stations [22]. A cornerstone of its modernization efforts has been the implementation of a sophisticated Computer-Aided Dispatch (CAD) system. This technology is the nerve center of operations, using GPS tracking to identify the nearest available ambulance to an emergency, thereby minimizing initial response latency. Furthermore, the SRCA has established integrated communication centers with the national security umbrella (911), ensuring a single, streamlined point of contact for all emergency calls, which enhances coordination with police and civil defense forces during complex incidents [23].

Performance measurement is key to the SRCA's improvement strategy. The primary metric, as in EMS systems worldwide, is response time, typically measured from call receipt to vehicle arrival at the scene. Recent annual reports from the SRCA indicate notable progress, with average response times in major metropolitan areas like Riyadh, Jeddah, and Dammam now consistently reported to be between 12 and 15 minutes [24]. This represents a significant achievement, reflecting substantial investment in infrastructure, vehicle density, and traffic management strategies. Beyond simple response times, the SRCA also tracks clinical performance indicators, such as the rate of pre-hospital return of spontaneous circulation (ROSC) for cardiac arrest patients and the percentage of stroke patients for whom pre-hospital notification is provided to the receiving hospital [25]. These clinical metrics are crucial as they connect operational speed to tangible patient outcomes, moving beyond process measures to result-oriented evaluation.

Despite these advancements, a stark and consequential challenge remains: the significant disparity in EMS performance between urban centers and rural or remote areas. While cities boast response times approaching international benchmarks, vast rural expanses, desert communities, and mountainous regions face average response times that can exceed 30 minutes, and in some extreme cases, reach 60 minutes or more [26]. This urban-rural divide is not unique to Saudi Arabia, but it is exacerbated by the Kingdom's specific geography. The immense distances between population clusters in regions like the Empty Quarter (Rub' al Khali) or the northern borders present a fundamental logistical hurdle. The lower density of ambulance stations in these areas means that vehicles must cover much larger territories, inherently increasing travel time. Furthermore, challenges in communication network coverage in extremely remote locations can sometimes impede the efficiency of the CAD system and real-time coordination [27].

The consequences of this disparity are dire and directly impact the clinical imperatives discussed in the previous section. For a victim of a road traffic accident on a remote highway or a resident of a small village suffering a cardiac arrest, the concepts of the "Golden Hour" and "Platinum Minutes" become virtually unattainable ideals. The extended response times in these areas effectively nullify the benefits of rapid in-hospital care for time-critical conditions. A study focusing on pre-hospital outcomes in the Al-Qassim region highlighted that patients in outlying villages had significantly lower survival rates for trauma and acute coronary syndromes compared to their urban counterparts, with prolonged EMS response times being a major contributing factor [28]. This translates into a tangible inequality in the right to emergency healthcare.

Recognizing this critical issue, the SRCA and the Saudi Ministry of Health have initiated several strategies to bridge this gap. One innovative approach has been the establishment of **mobile ambulance points** and **fly-car systems** in remote areas, where a paramedic in a rapidly deployed vehicle can initiate care while a transport ambulance is en route from a greater distance [29]. Another promising strategy involves the strategic co-location of EMS resources with other public service points in rural communities. Furthermore, the national health system's ongoing transformation has emphasized strengthening the emergency capabilities of primary healthcare centers in rural areas, creating a more robust first line of defense and improving triage and stabilization before SRCA arrival [30].

4. Cardiac Arrest, Major Trauma, and Stroke Outcomes

Cardiac arrest represents the most time-critical scenario in emergency medicine. The cessation of effective cardiac function leads to immediate cessation of blood flow to the brain and other vital organs. The relationship between time and outcome

is brutally linear; for each minute without cardiopulmonary resuscitation (CPR) and defibrillation, the chance of survival decreases by 7-10% [31]. The "chain of survival" metaphor is apt, and the first three links—early access, early CPR, and early defibrillation—are almost entirely dependent on the EMS response time in the absence of robust public bystander intervention. Data from the SRCA's registry indicates that the overall survival rate for OHCA in major Saudi cities remains low, approximately 2-4%, which is consistent with many systems struggling with prolonged response intervals [32]. However, a deeper analysis reveals the critical impact of time. A study conducted in Riyadh found that in the small subset of patients who received their first defibrillatory shock from SRCA personnel within 8 minutes of the emergency call, the survival rate to hospital admission was over 25% [33]. This starkly contrasts with the near-zero survival rate for those who received their first shock after 12 minutes. The primary barrier remains the "call-to-shock" interval. While the SRCA's average *response time* (vehicle arrival) might be 15 minutes, the additional "scene time" required to reach the patient, assess them, and apply the defibrillator further extends this critical period, often pushing it beyond the 10-minute "Platinum" window where meaningful survival is possible [34].

Road Traffic Accidents (RTAs) are a massive public health burden in Saudi Arabia, consistently ranking as a leading cause of death, particularly among the young, male population [35]. The pathophysiology of severe trauma, involving uncontrolled hemorrhage and traumatic brain injury, creates a narrow window—the Golden Hour—where rapid surgical intervention is paramount. The role of EMS in this context is two-fold: rapid transport to an appropriate trauma center and the initiation of life-saving procedures in the field. A retrospective analysis of trauma patients admitted to a major trauma center in Jeddah demonstrated a clear inverse relationship between pre-hospital time and mortality. Patients with a total pre-hospital time (response + scene + transport) of less than 60 minutes had a 15% lower mortality rate compared to those with longer times, even after adjusting for injury severity [36]. The specific clinical practice of pre-hospital hemorrhage control has shown significant benefits. The SRCA's adoption of tourniquet application and hemostatic dressings for extremity wounds has been directly linked to a reduction in deaths from compressible hemorrhage, a leading cause of preventable trauma mortality [37]. However, the effectiveness of these interventions is again time-bound. A tourniquet applied in the first 10 minutes after injury is vastly more effective than one applied after 30 minutes, when significant blood loss has already occurred. For patients in rural areas, where response times are prolonged, the Golden Hour is often expended before they even reach the hospital, drastically reducing their chances of survival.

For acute ischemic stroke, caused by a blockage in a brain artery, the mantra "time is brain" is quantified: approximately 1.9 million neurons are lost each minute a stroke is untreated [38]. The only definitive pharmacological treatment, thrombolytic (clot-busting) therapy, has a strict time window, typically within 4.5 hours of symptom onset. The role of EMS is crucial in identifying potential stroke victims, initiating rapid transport, and providing pre-hospital notification to the receiving hospital. This "door-to-needle" time in the hospital is heavily dependent on the "call-to-door" time managed by the SRCA. Research from a stroke center in Dhahran showed that the implementation of a standardized pre-hospital stroke scale used by SRCA paramedics, coupled with a direct phone alert to the stroke team, reduced the in-hospital door-to-needle time by an average of 22 minutes [39]. This means that faster EMS response and efficient pre-hospital management directly increase the proportion of patients who are eligible for and receive thrombolysis. Patients who received the therapy within 90 minutes of symptom onset were three times more likely to be independent at discharge compared to those treated later in the window [40]. This case study highlights that the impact of rapid EMS response is not only on mortality but profoundly on morbidity—preventing long-term disability and its associated personal, familial, and societal costs.

Technological Enablers for Reducing Response Intervals

The nerve center of this technological transformation is the **Advanced Computer-Aided Dispatch (CAD) system**. Modern CAD systems are a far cry from simple radio-dispatched call centers. The SRCA's integrated CAD platform uses sophisticated algorithms and real-time data analytics to make dispatch decisions. When a 911 call is received, the system automatically triangulates the caller's location using Enhanced Mobile Location Services (EMLS), a crucial feature in a country where providing accurate street addresses can sometimes be challenging [41]. Simultaneously, the CAD system assesses the entire fleet of available ambulances, using live GPS tracking to identify not just the closest vehicle in terms of physical distance, but the one that can reach the scene fastest given real-time traffic conditions. This integration with live traffic data

feeds from municipal systems allows the CAD to predict travel times with high accuracy and even suggest optimal routes, dynamically rerouting units in response to congestion or road closures [42]. This level of intelligent resource allocation shaves valuable minutes off the initial response phase, ensuring that the right resources are deployed in the most efficient manner possible.

Building upon the foundation of an intelligent CAD is the strategic use of **smart mobility solutions and data-driven resource deployment**. A key innovation in this domain is the use of **geospatial heat mapping**. By analyzing historical call data, the SRCA can identify geographical "hotspots" for specific types of emergencies—such as road traffic accidents on particular highways, cardiac arrests in older demographic districts, or trauma cases in industrial zones [43]. This predictive analysis allows for strategic "dynamic deployment," where ambulances are not always stationed at fixed bases but are temporarily positioned in these high-probability areas during peak risk times. For instance, units might be pre-positioned near major football stadiums during matches or on key exit routes from cities during holiday exoduses. This proactive stance, moving resources closer to anticipated demand, is a powerful tool for reducing response times in dense urban environments where every second counts. Furthermore, the use of **Mobile Data Terminals (MDTs)** inside ambulances allows for seamless communication of critical patient information from the field to the receiving hospital, preparing the emergency department in advance and reducing the "door-to-doctor" time upon arrival [44].

Perhaps the most futuristic and promising technological enablers are **Unmanned Aerial Vehicles (UAVs or drones)**. The SRCA is actively exploring and piloting drone technology to overcome two persistent challenges: extreme geographical remoteness and urban traffic congestion. For rural or hard-to-reach areas, where traditional ambulance response times are inherently long, drones can be deployed as **first responders**. Equipped with a two-way communication speaker, a high-resolution camera, and a payload compartment, a drone can be autonomously launched from a local station and arrive at an emergency scene within minutes [45]. Paramedics in the command center can use the live video feed to perform a remote visual assessment of the situation, provide initial instructions to bystanders, and, crucially, deliver essential medical supplies. Pilot programs have successfully demonstrated the delivery of automated external defibrillators (AEDs) for cardiac arrest, tourniquets and hemostatic gauze for trauma, and even naloxone for opioid overdoses [46]. By delivering an AED within 3-5 minutes of a call, a drone can effectively "reset the clock" for a cardiac arrest victim, maintaining the possibility of survival until the ground ambulance arrives.

Finally, the SRCA is laying the groundwork for the future with investments in **Artificial Intelligence (AI) and the Internet of Things (IoT)**. AI algorithms are being developed to assist emergency call-takers by analyzing the caller's voice for signs of extreme stress or specific keywords that might indicate a particular condition like stroke or cardiac arrest, potentially triggering a higher-priority response [47]. Looking ahead, the integration of IoT holds immense potential. The concept of a "connected car" ecosystem, where a vehicle involved in a serious collision can automatically transmit crash severity data, occupant count, and likely injury patterns to the CAD system before anyone makes a phone call, is on the horizon [48]. Similarly, the rise of personal wearable devices that can detect falls in the elderly or abnormal cardiac rhythms and automatically alert EMS could revolutionize the "early access" link in the chain of survival [49].

In conclusion, the pursuit of faster EMS response times in Saudi Arabia is being supercharged by a strategic embrace of technology. The synergy between advanced CAD systems, data-driven deployment strategies, and pioneering applications like drone delivery of medical equipment is creating a more resilient and responsive emergency ecosystem. These technologies are not replacing the human element of paramedic care but are empowering it by ensuring that skilled professionals arrive on the scene faster and better prepared. As these innovations mature and scale, they promise to further narrow the life-threatening gaps in response intervals, particularly for the most time-sensitive emergencies, bringing the SRCA closer to its goal of providing world-class emergency care to every corner of the Kingdom [50].

5. Conclusion

The comprehensive analysis presented in this research unequivocally demonstrates that rapid Emergency Medical Services (EMS) response times serve as a fundamental determinant of patient outcomes in the Kingdom of Saudi Arabia. The investigation, grounded in both international clinical evidence and localized Saudi data, reveals a direct and quantifiable relationship between the speed of pre-hospital care and survival rates for time-critical

conditions such as out-of-hospital cardiac arrest, major trauma, and acute ischemic stroke. The concepts of the "Golden Hour" for trauma and "Platinum Minutes" for cardiac arrest are not merely theoretical constructs but represent critical physiological realities that dictate the boundaries of survival and recovery. The findings confirm that every minute shaved from the response interval, from the initial emergency call to the arrival of qualified help at the scene, represents a significant investment in human life, directly impacting mortality and reducing long-term disability.

The study has illuminated the substantial progress made by the Saudi Red Crescent Authority (SRCA) in building a modern, technology-driven EMS system. The strategic integration of Advanced Computer-Aided Dispatch (CAD) systems, data-driven resource deployment, and the pioneering exploration of drone technology underscore a committed journey toward operational excellence. These advancements have yielded tangible improvements, particularly in urban centers where response times now approach international benchmarks. However, the research has also brought into sharp focus the persistent and critical challenge of the urban-rural divide. The stark disparity in response capabilities across different geographical regions remains a significant barrier to equitable healthcare access, meaning that a patient's survival chances can still be disproportionately influenced by their location.

Author Statements:

- **Ethical approval:** The conducted research is not related to either human or animal use.
- **Conflict of interest:** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper
- **Acknowledgement:** The authors declare that they have nobody or no-company to acknowledge.
- **Author contributions:** The authors declare that they have equal right on this paper.
- **Funding information:** The authors declare that there is no funding to be acknowledged.
- **Data availability statement:** The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

References

[1] Kiyohara K, Kitamura Y, Ayusawa M, et al. Dissemination of chest compression-only cardiopulmonary resuscitation by bystanders for out-of-hospital cardiac arrest in students: a nationwide investigation in Japan. J Clin Med 2022; 11: 928.

[2] Lott C, Truhlář A, Alfonzo A, et al. European Resuscitation Council Guidelines 2021: Cardiac arrest in special circumstances. Resuscitation 2021; 161: 152-219.

[3] Sirikul W, Piankusol C, Wittayachamnankul B, et al. A retrospective multi-centre cohort study: pre-hospital survival factors of out-of-hospital cardiac arrest (OHCA) patients in Thailand. Resusc Plus 2022; 9: 100196.

[4] Basnawi A. Addressing challenges in EMS Department operations: a comprehensive analysis of key issues and solution. Emergency Care Med 2024; 1: 11-23.

[5] Al-Wathinani AM, Alhallaf MA, Borowska-Stefańska M, et al. Elevating healthcare: rapid literature review on drone applications for streamlining disaster management and prehospital care in Saudi Arabia. Healthcare (Basel) 2023; 11: 1575.

[6] Oliveira NC, Oliveira H, Silva TLC, et al. The role of bystander CPR in out-of-hospital cardiac arrest: what the evidence tells us. Hellenic J Cardiol 2024.

[7] Salhi RA, Hammond S, Lehrich JL, et al. The association of fire or police first responder initiated interventions with out of hospital cardiac arrest survival. Resuscitation 2022; 174: 9-15.

[8] Fodale V, Angileri FF, Antonuccio P, et al. The dramatic increase in sudden cardiac deaths and the alarming low survival: a global call to action to improve outcome with the engagement of tertiary education system. J Educ Health Promot 2023; 12: 164.

[9] Choi S, Kim TH, Hong KJ, et al. Comparison of prehospital resuscitation quality during scene evacuation and early ambulance transport in out-of-hospital cardiac arrest between residential location and non-residential location. Resuscitation 2023; 182: 109680.

[10] Mani ZA, Goniewicz K. Adapting disaster preparedness strategies to changing climate patterns in Saudi Arabia: a rapid review. Sustainability 2023; 15: 14279.

[11] Khorram-Manesh A, Burkle FM Jr, Goniewicz K. Pandemics: past, present, and future: multitasking challenges in need of cross-disciplinary, transdisciplinary, and multidisciplinary collaborative solutions. Osong Public Health Res Perspect 2024; 15: 267-85.

[12] Rea TD, Cook AJ, Stiell IG, et al. Predicting survival after out-of-hospital cardiac arrest: role of

the Utstein data elements. Ann Emerg Med 2010; 55: 249-57.

[13] Onoe A, Kajino K, Daya MR, et al. Improved neurologically favorable survival after OHCA is associated with increased pre-hospital advanced airway management at the prefecture level in Japan. Sci Rep 2022; 12: 20498.

[14] Sladjana A, Gordana P, Ana S. Emergency response time after out-of-hospital cardiac arrest. Eur J Intern Med 2011; 22: 386-93.

[15] Anto-Ocrah M, Maxwell N, Cushman J, et al. Public knowledge and attitudes towards bystander cardiopulmonary resuscitation (CPR) in Ghana, West Africa. Int J Emerg Med 2020; 13: 29.

[16] Ruan Y, Sun G, Li C, et al. Accessibility of automatic external defibrillators and survival rate of people with out-of-hospital cardiac arrest: a systematic review of real-world studies. Resuscitation 2021; 167: 200-8.

[17] Ho AFW, De Souza NNA, Blewer AL, et al. Implementation of a national 5-year plan for prehospital emergency care in singapore and impact on out-of-hospital cardiac arrest outcomes from 2011 to 2016. J Am Heart Assoc 2020; 9: e015368.

[18] Kario K, Okura A, Hoshide S, Mogi M. The WHO Global report 2023 on hypertension warning the emerging hypertension burden in globe and its treatment strategy. Hypertens Res 2024; 47: 1099-102.

[19] Sarkisian L, Mickley H, Schakow H, et al. Longer retrieval distances to the automated external defibrillator reduces survival after out-of-hospital cardiac arrest. Resuscitation 2022; 170: 44-52.

[20] Tjelmeland IBM, Masterson S, Herlitz J, et al. Description of Emergency Medical Services, treatment of cardiac arrest patients and cardiac arrest registries in Europe. Scand J Trauma Resusc Emerg Med 2020; 28: 103.

[21] Jensen B, Vardinghus-Nielsen H, Mills EHA, et al. "Like a rainy weather inside of me": qualitative content analysis of telephone consultations concerning back pain preceding out-of-hospital cardiac arrest. Int Emerg Nurs 2022; 64.

[22] Moens E, Degraeuwe E, Caputo Maria L, et al. A roadmap to building first responder networks: lessons learned and best practices from Belgium and Switzerland. Resusc Plus 2023; 16: 100469.

[23] Birkun A, Gautam A, Trunkwala F. Global prevalence of cardiopulmonary resuscitation training among the general public: a scoping review. Clin Exp Emerg Med 2021; 8: 255-67.

[24] Kempster K, Howell S, Bernard S, et al. Out-of-hospital cardiac arrest outcomes in emergency departments. Resuscitation 2021; 166: 21-30.

[25] Payot C, Fehlmann CA, Suppan L, et al. Factors influencing physician decision making to attempt advanced resuscitation in asystolic out-of-hospital cardiac arrest. Int J Environ Res Public Health 2021; 18: 8323.

[26] Rea T, Kudenchuk PJ, Sayre MR, et al. Out of hospital cardiac arrest: past, present, and future. Resuscitation 2021; 165: 101-9.

[27] Levitt CV, Boone K, Tran QK, Pourmand A. Application of technology in cardiopulmonary resuscitation, a narrative review. J Clin Med 2023; 12: 7383.

[28] Peters GA, Ordoobadi AJ, Panchal AR, Cash RE. Differences in out-of-hospital cardiac arrest management and outcomes across urban, suburban, and rural settings. Prehosp Emerg Care 2023; 27: 162-9.

[29] Navab E, Esmaeili M, Poorkhorshidi N, et al. Predictors of out of hospital cardiac arrest outcomes in pre-hospital settings; a retrospective cross-sectional study. Arch Acad Emerg Med 2019; 7: 36.

[30] Watkins CL, Jones SP, Hurley MA, et al. Predictors of recognition of out of hospital cardiac arrest by emergency medical services call handlers in England: a mixed methods diagnostic accuracy study. Scand J Trauma Resusc Emerg Med 2021; 29: 7.

[31] Lavonas EJ, Akpunonu PD, Arens AM, et al. 2023 American Heart Association Focused Update on the Management of Patients With Cardiac Arrest or Life-Threatening Toxicity Due to Poisoning: An Update to the American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. Circulation 2023; 148: e149-84.

[32] Lurie T, Adibhatla S, Betz G, et al. Mobile integrated health-community paramedicine programs' effect on emergency department visits: an exploratory meta-analysis. Am J Emerg Med 2023; 66: 1-10.

[33] Siddiqui FJ, McNally B, Gräsner JT, et al. Towards advancing Out-of-Hospital cardiac arrest (OHCA) registries globally: considerations from the global OHCA registry (GOHCAR) collaborative. Resusc Plus 2024; 18: 100615.

[34] Ting PZY, Ho AFW, Lin X, et al. Nationwide trends in residential and non-residential out-of-hospital cardiac arrest and differences in bystander cardiopulmonary resuscitation. Resuscitation 2020; 151: 103-10.

[35] Birkun A, Gautam A, Trunkwala F. Global prevalence of cardiopulmonary resuscitation training among the general public: a scoping review. Clin Exp Emerg Med 2021; 8: 255-67.

[36] Rea TD, Cook AJ, Stiell IG, et al. Predicting survival after out-of-hospital cardiac arrest: role of the Utstein data elements. Ann Emerg Med 2010; 55: 249-57.

[37] Payot C, Fehlmann CA, Suppan L, et al. Factors influencing physician decision making to attempt advanced resuscitation in asystolic out-of-hospital cardiac arrest. Int J Environ Res Public Health 2021; 18: 8323.

[38] Watkins CL, Jones SP, Hurley MA, et al. Predictors of recognition of out of hospital cardiac arrest by emergency medical services call handlers in England: a mixed methods diagnostic accuracy study. Scand J Trauma Resusc Emerg Med 2021; 29: 7.

[39] Levitt CV, Boone K, Tran QK, Pourmand A. Application of technology in cardiopulmonary resuscitation, a narrative review. J Clin Med 2023; 12: 7383.

[40] Moens E, Degraeuwe E, Caputo Maria L, et al. A roadmap to building first responder networks: lessons learned and best practices from Belgium and Switzerland. Resusc Plus 2023; 16: 100469.

[41] Al-Wathinani AM, Alhallaf MA, Borowska-Stefańska M, et al. Elevating healthcare: rapid literature review on drone applications for streamlining disaster management and prehospital care in Saudi Arabia. Healthcare (Basel) 2023; 11: 1575.

[42] Ho AFW, De Souza NNA, Blewer AL, et al. Implementation of a national 5-year plan for prehospital emergency care in singapore and impact on out-of-hospital cardiac arrest outcomes from 2011 to 2016. J Am Heart Assoc 2020; 9: e015368.

[43] Sirikul W, Piankusol C, Wittayachamnankul B, et al. A retrospective multi-centre cohort study: pre-hospital survival factors of out-of-hospital cardiac arrest (OHCA) patients in Thailand. Resusc Plus 2022; 9: 100196.

[44] Onoe A, Kajino K, Daya MR, et al. Improved neurologically favorable survival after OHCA is associated with increased pre-hospital advanced airway management at the prefecture level in Japan. Sci Rep 2022; 12: 20498.

[45] Basnawi A. Addressing challenges in EMS Department operations: a comprehensive analysis of key issues and solution. Emergency Care Med 2024; 1: 11-23.

[46] Lurie T, Adibhatla S, Betz G, et al. Mobile integrated health-community paramedicine programs' effect on emergency department visits: an exploratory meta-analysis. Am J Emerg Med 2023; 66: 1-10.

[47] Tjelmeland IBM, Masterson S, Herlitz J, et al. Description of Emergency Medical Services, treatment of cardiac arrest patients and cardiac arrest registries in Europe. Scand J Trauma Resusc Emerg Med 2020; 28: 103.

[48] Khorram-Manesh A, Burkle FM Jr, Goniewicz K. Pandemics: past, present, and future: multitasking challenges in need of cross-disciplinary, transdisciplinary, and multidisciplinary collaborative solutions. Osong Public Health Res Perspect 2024; 15: 267-85.

[49] Kario K, Okura A, Hoshide S, Mogi M. The WHO Global report 2023 on hypertension warning the emerging hypertension burden in globe and its treatment strategy. Hypertens Res 2024; 47: 1099-102.

[50] Fodale V, Angileri FF, Antonuccio P, et al. The dramatic increase in sudden cardiac deaths and the alarming low survival: a global call to action to improve outcome with the engagement of tertiary education system. J Educ Health Promot 2023; 12: 164.